

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



✓



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07H 19/04, 21/00</p>	<p>A2</p>	<p>(11) International Publication Number: WO 97/47636 (43) International Publication Date: 18 December 1997 (18.12.97)</p>
<p>(21) International Application Number: PCT/GB97/01490 (22) International Filing Date: 3 June 1997 (03.06.97) (30) Priority Data: 9612600.8 13 June 1996 (13.06.96) GB (71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): COLLINGWOOD, Stephen, Paul [GB/GB]; 14 Fowler Close, Maidenbower, Crawley, West Sussex RH10 7UN (GB). MOSER, Heinz, Ernst [CH/GB]; 32 Hazel Close, Southwater, Horsham, West Sussex RH13 7GN (GB). ALTMANN, Karl-Heinz [DE/CH]; Tannenweg 5, CH-4153 Reinach (CH). DOUGLAS, Mark, Edward [GB/GB]; 71 High Street, Macclesfield, Cheshire SK11 7QQ (GB). (74) Agent: MARSH, Bernard, Andrew; Novartis UK Limited, Patent and Trademark Dept., Central Research, Hulley Road, Macclesfield SK10 2NX (GB).</p>		<p>(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.</p>

(54) Title: INTERMEDIATES FOR OLIGONUCLEOTIDE SYNTHESIS

(57) Abstract

A compound of formula (I) where B¹ is a radical of a nucleoside base, R¹ is hydrogen or a hydroxy-protecting group, R² is hydrogen, hydroxy or a 2' nucleoside-modifying atom or group, R³ is an unsubstituted or substituted C₁ to C₁₀ alkyl, C₂ to C₁₀ alkenyl, C₄ to C₁₀ cycloalkylalkyl, C₆ to C₁₀ aryl or C₇ to C₁₃ aralkyl group, and Z is halogen or a group of formula (II) where R⁴ and R⁵ are each independently an unsubstituted or substituted C₁ to C₁₀ alkyl, C₂ to C₁₀ alkenyl, C₄ to C₁₀ cycloalkylalkyl, C₆ to C₁₀ aryl or C₇ to C₁₃ aralkyl group, or R⁴ is said group and R⁵ is hydrogen, or R⁴ and R⁵ together with the nitrogen atom to which they are attached denote a five- to thirteen-membered heterocyclic ring, or Z is a group of formula (III): Nuc-O-, where Nuc is the residue of a natural or synthetic nucleoside or oligonucleotide after removal of a 5'-hydroxyl group therefrom attached through a 5' methylene thereof to the indicated oxygen atom.



FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		


INTERMEDIATES FOR OLIGONUCLEOTIDE SYNTHESIS

This invention relates to chemical compounds which are intermediates for oligonucleotides and the use of such compounds in the preparation of oligonucleotides.


Accordingly, the present invention provides, in one aspect, a compound of formula



where B^1 is a radical of a nucleoside base, R^1 is hydrogen or a hydroxy-protecting group, R^2 is hydrogen, hydroxy or a 2' nucleoside-modifying atom or group, R^3 is an unsubstituted or substituted C_1 to C_{10} alkyl, C_2 to C_{10} alkenyl, C_4 to C_{10} cycloalkylalkyl, C_6 to C_{10} aryl or C_7 to C_{13} aralkyl group, and Z is halogen or a group of formula



where R^4 and R^5 are each independently an unsubstituted or substituted C_1 to C_{10} alkyl, C_2 to C_{10} alkenyl, C_4 to C_{10} cycloalkylalkyl, C_6 to C_{10} aryl or C_7 to C_{13} aralkyl group, or R^4 is said group and R^5 is hydrogen, or R^4 and R^5 together with the nitrogen atom to which they are attached denote a five- to thirteen- membered heterocyclic ring; or Z is a group of formula



where Nuc is the residue of a natural or synthetic nucleoside or oligonucleotide after removal of a 5' hydroxyl group therefrom attached through a 5' methylene thereof to the indicated oxygen atom.

When R^3 , R^4 or R^5 denote a substituted alkyl, alkenyl, cycloalkylalkyl, aryl or aralkyl group, the group is preferably substituted by halogen, hydroxy, C_1 - C_4 alkoxy, cyano, nitro or tri(C_1 - C_{15} hydrocarbyl)silyl or, in the case of R^3 , by C_1 - C_4 alkylsulphonyl or C_6 - C_{10} arylsulphonyl.

When a tri(C_1 - C_{15} hydrocarbyl)silyl group is present in compounds of formula I, it may be, for example, trialkylsilyl such as trimethylsilyl, triethylsilyl, tri-n-propylsilyl, tri-isopropylsilyl, tri-n-butylsilyl, tri-isobutylsilyl, isopropyl dimethylsilyl, tert-butyl dimethylsilyl or 1,1,2,2-tetramethylethyl dimethylsilyl (thexyl dimethylsilyl), triarylsilyl such as triphenylsilyl, triarylalkylsilyl such as tribenzylsilyl, aryl dialkylsilyl such as phenyl dimethylsilyl, phenyl diethylsilyl, phenyl diisopropylsilyl or phenyl di-tert-butylsilyl, or alkyl diarylsilyl such as methyl diphenylsilyl, isopropyl diphenylsilyl or tert-butyl diphenylsilyl. Where R^3 , R^4 or R^5 has a tri(C_1 - C_{15} hydrocarbyl)silyl substituent, it is preferably tri(C_1 - C_4 hydrocarbyl)silyl, especially trimethylsilyl. R^1 as tri(C_1 - C_{15} hydrocarbyl)silyl is preferably C_1 - C_6 alkyl di (C_6 - C_8 aryl)silyl, especially tert-butyl diphenylsilyl, or branched C_2 - C_{10} alkyl di (C_1 - C_4 alkyl)silyl, especially hexyl dimethylsilyl.

R^1 may be any hydroxy-protecting group capable of protecting a 5'-hydroxyl group against undesired reaction. Such groups are well known and include C_1 to C_{10} aliphatic, e.g. alkyl groups; C_3 to C_8 cycloaliphatic, e.g. cycloalkyl, groups; C_6 to C_{10} aromatic, e.g. aryl, groups; C_7 to C_{40} araliphatic, e.g. aralkyl or C_1 to C_4 -alkoxy-substituted aralkyl groups; groups of formula -COR or -SO₂R where R is a C_1 to C_{10} aliphatic group, a C_3 to C_8 cycloaliphatic group, a C_6 to C_{10} aromatic group or a C_7 to C_{40} araliphatic group; and tri(C_1 - C_{15} hydrocarbyl)silyl groups. Preferably R^1 is hydrogen or a 5'-protecting group conventionally used in oligonucleotide synthesis, more preferably hydrogen, a C_7 to C_{40} araliphatic group or a tri(C_1 - C_{15} hydrocarbyl)silyl group, especially an unsubstituted or substituted triphenylmethyl group such as methoxytrityl, dimethoxytrityl or tris tert-butyltrityl or a C_1 - C_6

alkyldi(C₆-C₈ aryl)silyl group such as tert-butyldiphenylsilyl, particularly dimethoxytrityl or tert-butyldiphenylsilyl.

R², the 2' nucleoside-modifying atom or group, i.e. an atom or group which may be attached to the 2' position of a nucleoside in place of a hydrogen atom or hydroxy group to effect a modification, may be a halogen atom such as a fluorine, chlorine or bromine atom; C₁ to C₁₀ unsubstituted alkyl, such as methyl, trifluoromethyl, ethyl, propyl, butyl, pentyl, hexyl, octyl or decyl; C₆ to C₁₀ aryl such as phenyl, tolyl or xylyl; C₇ to C₁₃ aralkyl such as benzyl; amino, C₁ to C₁₀ alkyl amino such as methylamino, ethylamino or octylamino; C₁ to C₁₀ alkylthio such as methylthio, ethylthio or octylthio; azide; nitrate; nitrite; cyanide; cyanate; methanesulphonate; C₁ to C₁₀ aminoalkylamino; a group of formula -OR⁶ or -OCOR⁶ where R⁶ is a C₁ to C₃₀ organic group; substituted silyl; an RNA cleaving group; a cholesteryl group; a conjugate; a reporter group; an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide.

Preferably R² is hydrogen, halogen, hydroxy or a group of formula -OR⁶ or -OCOR⁶ where R⁶ is a C₁ to C₁₀ alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, n-heptyl, n-octyl, 2-ethylhexyl, n-nonyl or n-decyl, C₂ to C₁₀ alkenyl group such as vinyl, allyl, methallyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl or decenyl, a C₃ to C₈ cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methylcyclopentyl, cyclohexyl, methylcyclohexyl, cycloheptyl or cyclooctyl, a C₄ to C₁₀ cycloalkylalkyl group such as cyclopropylmethyl, cyclobutylmethyl, cyclohexylmethyl, methylcyclohexylmethyl, cyclohexyl ethyl or dimethyl cyclohexylethyl, a C₆ to C₁₀ aryl group such as phenyl, ortho-, meta- or para-tolyl, ortho-, meta- or para-xylyl or naphthyl or a C₇ to C₂₅ aralkyl group such as benzyl, phenylethyl, diphenylmethyl or triphenylmethyl, any of which groups may be unsubstituted or substituted, for example by halogen, hydroxy, C₁ to C₄ alkoxy or cyano, or, where R² is of formula -OR⁶, R⁶ is a tri(C₁-C₁₅ hydrocarbyl)silyl group.

In more preferred embodiments, R^2 is hydrogen, hydroxy, fluorine, a group of formula $-OR^6$ where R^6 is a C_1 to C_4 alkyl group, a C_2 to C_4 alkenyl group, a C_7 to C_{13} aralkyl group or a group of formula $-(R^7-O)_n-R^8$ where R^7 is C_1 to C_4 alkylene, R^8 is C_1 to C_4 alkyl and n is 1 to 8, or R^2 is a group of formula $-OCOR^6$ where R^6 is C_1 to C_4 alkyl or C_6 to C_8 aryl. In especially preferred embodiments, R^2 is hydrogen, hydroxy, fluorine, methoxy, ethoxy, allyloxy, benzyloxy, methoxyethoxy, $-O-(CH_2-CH_2-O)_3-CH_3$, acetoxy or benzoyloxy.

R^3 , R^4 or R^5 as C_1 to C_{10} alkyl may be, for example a straight or branched group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, n-heptyl, n-octyl, 2-ethylhexyl, n-nonyl or n-decyl. R^3 , R^4 or R^5 as C_2 to C_{10} alkenyl may be, for example, vinyl, allyl, methallyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl or decenyl. R^3 , R^4 or R^5 as C_4 to C_{10} cycloalkylalkyl may be, for example, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclopentylethyl, methylcyclopentylmethyl, cyclohexylmethyl, cyclohexylethyl, methylcyclohexylmethyl, methylcyclohexylethyl, dimethylcyclohexylmethyl or dimethylcyclohexylethyl. R^3 , R^4 or R^5 as C_6 to C_{10} aryl may be, for example, phenyl, o-tolyl, m-tolyl, p-tolyl, o-xylyl, m-xylyl or p-xylyl. R^3 , R^4 or R^5 as C_7 to C_{13} aralkyl may be, for example, benzyl, phenylethyl or diphenylmethyl. Any of the above groups as R^3 , R^4 or R^5 may be unsubstituted or substituted as hereinbefore described.

When R^4 and R^5 together with the attached nitrogen atom denote a heterocyclic ring, this ring may be, for example, a five-membered ring such as a pyrrolidine ring, a six-membered ring such as a piperidine, 2,6-dimethylpiperidine, N-methylpiperazine, morpholine or thiomorpholine ring, a seven-membered ring such as a perhydroazepine ring, an eight-membered ring such as a perhydroazocine ring, a nine-membered ring such as perhydroazonine ring, a ten-membered ring such as a perhydroazecine ring, an eleven-membered ring such as an azacycloundecane ring, a twelve-membered ring such as an azacyclododecane ring or a thirteen-membered ring such as an azacyclotridecane ring.

R^3 is preferably unsubstituted or substituted C_1 to C_4 alkyl, C_3 to C_5 alkenyl, C_6 to C_8 cycloalkylalkyl or C_7 to C_9 aralkyl, especially methyl, ethyl, 2-cyanoethyl, 2,2,2-trichloroethyl, allyl, benzyl or 2-(trimethylsilyl)ethyl.

When Z in formula I is a group of formula II, in preferred embodiments R^4 and R^5 independently are unsubstituted or substituted C_1 to C_4 alkyl, C_3 to C_5 alkenyl, C_6 to C_8 cycloalkylalkyl or C_7 to C_9 aralkyl, or R^4 is C_1 to C_4 alkyl, preferably branched C_1 to C_4 alkyl such as isopropyl or tert-butyl, or C_7 to C_9 aralkyl such as benzyl, and R^5 is hydrogen, or R^4 and R^5 together with the attached nitrogen atom denote a five- or six- membered heterocyclic ring having one or two heteroatoms in the ring. In especially preferred embodiments, R^4 and R^5 are independently methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, allyl or benzyl, or R^4 and R^5 together with the attached nitrogen atom denote a pyrrolidine, piperidine or morpholine ring.


When Z in formula I is a group of formula III, the residue Nuc may be the residue, after removal of a 5'-hydroxyl group, of any natural or synthetic nucleoside, for example adenosine, cytidine, guanosine, thymidine, uridine, 2-aminoadenosine, 6-N-benzyladenosine, 5-N-methylcytidine, 4-N-benzoylcytidine, 5-propynylcytidine, 2-N-isobutyrylguanosine, N-benzyloxymethylthymidine or 5-fluorouracil, or an oligonucleotide derived therefrom which may be a dinucleotide, trinucleotide or higher oligomer, and which may have natural, i.e. phosphodiester, internucleoside linkages or synthetic analogues thereof such as phosphorothioate or methylphosphonate linkages, or P (III) precursors of such internucleoside linkages.

B^1 in formula I may be a radical of a base found in naturally occurring nucleosides, which may be unsubstituted or substituted, for example on a nitrogen atom by a protecting group, e.g. an acyl group, which may be aliphatic such as acetyl, or aromatic such as benzoyl or nitrobenzoyl, or by an aralkyloxyalkyl protecting group such as benzyloxymethyl, or synthetic analogues of such bases. Other types of protecting group conventionally used in nucleotide chemistry which may be present on a nitrogen atom in B^1 include amidine groups, such as N, N-dimethyl formamidine, N,N-dimethylacetamidine and N-methylpyrrolidine amidine groups, and alkoxy carbonyl groups such as tert-butoxy carbonyl which convert an amino group into a carbamate group.

Thus B¹ may be a radical of a purine or pyrimidine base such as adenine, guanine, cytosine, thymine or uracil, or an analogue of these bases such as 2-aminoadenine, 6-hydroxypurine, 5-methylcytosine, 5-propynylcytosine, 5-fluorouracil, 5-propynyluracil or dihydrouracil, or such a radical substituted on a nitrogen atom, for example by a protecting group as hereinbefore described.

In certain preferred embodiments where Z is a group of formula III, Nuc is a residue of a nucleoside or oligonucleotide as hereinbefore described attached at a terminal 3' position to a solid support such as a support used in solid phase oligonucleotide synthesis, for example to an inert silica-based support, such as controlled pore glass (CPG), containing long chain alkylamino groups, by a linker such as a succinyl group. The solid support may also contain groups to act as 3' terminal modifying groups for an oligonucleotide to be synthesised on the support.


Compounds of the invention may be in the form of one of the possible isomers, for example as a diastereoisomer, an optical isomer, a racemate or a mixture thereof. Preferred isomers of compounds of formula I are those of formula




where B¹, R¹, R² and R³ are as hereinbefore defined.

Specific especially preferred compounds of formula I are those hereinafter described in the Examples.

Compounds of formula I where Z is a group of formula II may be prepared by reacting a compound of formula I where Z is halogen, i.e. a compound of formula




where B¹, R¹, R² and R³ are as hereinbefore defined and Y is halogen, with a compound of formula



where R⁴ and R⁵ are as hereinbefore defined. The reaction may be carried out in an organic solvent, for example a halogenated hydrocarbon such as chloroform, in the presence of a tertiary nitrogen base such as pyridine, and at a temperature from -78°C to 50°C, preferably from -30°C to 25°C.

Compounds of formula V may be prepared by non-oxidative halogenation of a compound of formula



where B¹, R¹, R² and R³ are as hereinbefore defined. Non-oxidative halogenation may be carried out by reacting the compound of formula VII with a non-oxidative halogenating agent, for example a halophosphorane such as triphenyldichlorophosphorane or dichloro tris (2,4,6-tribromophenoxy)phosphorane in the presence of a base, preferably a tertiary

nitrogen base such as pyridine, in an organic solvent, which may be pyridine but is preferably a halohydrocarbon such as chloroform, at a temperature from -20°C to 60°C, preferably from 0°C to 50°C.

Compounds of formula VI are amines which are commercially available or may be prepared by known procedures.


Compounds of formula VII and their preparation are described in WO 96/08503.

Compounds of formula I where Z is a group of formula III may be prepared by reacting a compound of formula I where R¹ is a hydroxy-protecting group and Z is a group of formula II with a natural or synthetic nucleoside or oligonucleotide having a free 5' hydroxyl group (including oligonucleotides having one or more P(III) precursor internucleoside linkages). The reaction may be carried out in solution or with the nucleoside or oligonucleotide attached to a solid carrier, as hereinbefore described. It has been found that the reaction may be carried out under conditions, and using procedures, known for coupling reactions between a nucleoside having a 3' phosphoramidite group and a nucleoside or oligonucleotide having a free 5'hydroxyl group in conventional oligonucleotide synthesis - see, for example 'Oligonucleotides and Analogues, A Practical Approach' ed. F. Eckstein, IRL Press 1991.

Thus the compound of formula I where Z is a group of formula II may be reacted with the nucleoside or oligonucleotide in the presence of an amine-protonating coupling catalyst (activating agent) such as tetrazole or 5-(4-nitrophenyl)tetrazole to give a compound of formula I where Z is a group of formula III. The reaction may be carried out at -20 to 50°C, preferably at room temperature.

When the compound of formula I where Z is a group of formula III is reacted with a nucleoside or oligonucleotide attached to a solid support, the product is a compound of formula I in which Z is a group of formula III where Nuc is a residue of a nucleoside or oligonucleotide attached to a solid support.

Compounds of formula I where Z is a group of formula III are intermediates for the preparation of oligonucleotides having more nucleoside units than an oligonucleotide residue in Nuc. Accordingly, the present invention also provides a method of preparing an oligonucleotide of formula




where B^1 , R^1 , R^3 and Nuc are as hereinbefore defined and X is oxygen or sulphur, which comprises subjecting a compound of formula I where Z is a group of formula III to oxidation to form an oligonucleotide of formula VIII where X is oxygen or to sulphurisation to give an oligonucleotide of formula VIII where X is sulphur.


It has been found that oxidation or sulphurisation of compounds of formula I where Z is a group of formula III may be effected by methods used for oxidation or sulphurisation respectively of phosphite internucleoside linkages. Thus oxidation of compounds of formula I where Z is a group of formula III may be effected by treatment with iodine and water, or with a hydroperoxide such as tert-butyl hydroperoxide, for example using conditions and procedures known for oxidation of phosphite internucleoside linkages in oligonucleotide synthesis. Sulphurisation of compounds of formula I where Z is a group of formula III may be effected by treatment with sulphur in the presence of a tertiary amine in an organic solvent, usually carbon disulphide, by treatment with [3H] 1,2-benzodithiol-3-one-1,1-dioxide (Beaucage reagent) or by treatment with tetraethylthiuram, for example using procedures

known for sulphurisation of phosphite internucleoside linkages. Both oxidation and sulphurisation are generally carried out at room temperature.


Accordingly, the invention provides in another aspect a method of preparing an oligonucleotide which comprises subjecting a nucleoside having a protected 5' hydroxy group and, at the 3' position, a group of formula



where R^3 , R^4 and R^5 are as hereinbefore defined, to a nucleoside coupling reaction with a natural or synthetic nucleoside or oligonucleotide having a free 5' hydroxy group, to form an oligonucleotide precursor having an internucleoside linkage of formula



where R^3 is as hereinbefore defined, and converting the precursor into an oligonucleotide having an internucleoside linkage of formula



where R^3 is as hereinbefore defined and X is oxygen or sulphur by oxidising the precursor to give an oligonucleotide having an internucleoside linkage of formula XI where X is

oxygen or sulphurising the precursor to give an oligonucleotide having an internucleoside linkage of formula XI where X is sulphur.

The nucleoside having at the 3' position a group of formula IX, the nucleoside or oligonucleotide having a free 5' hydroxy group and the methods of carrying out the reactions may be as hereinbefore described.

Oligonucleotides of formula VIII where Nuc is a residue of a nucleoside or oligonucleotide attached to a solid support may be treated to remove the 5' protecting group R¹ and the resulting 5' hydroxy-terminated oligonucleotide subjected to successive coupling reactions with a natural or synthetic nucleoside or oligonucleotide having a protected 5' hydroxyl group and, at the 3' position, a group reactive with, or activatable to be reactive with, the free 5' hydroxy group on the deprotected oligonucleotide of formula VIII attached to the solid support, until an oligonucleotide of the desired length is obtained. Thus the oligonucleotide of formula VIII may be coupled with a nucleoside or oligonucleotide having a 3' phosphoramidite, H-phosphonate or phosphodiester group and a protected 5' hydroxyl group, or with a compound of formula I where Z is a group of formula II, to give a chain-extended oligonucleotide which may in turn be further chain extended by further such alternative reactions until an oligonucleotide of the desired length is obtained. Where a nucleoside or oligonucleotide having a 3' phosphoramidite, H-phosphonate or phosphodiester group is used, the coupling reactions may be carried out using procedures known in oligonucleotide synthesis. Where a compound of formula I where Z is a group of formula II is used, the coupling reaction may be carried out as hereinbefore described. Thus, where a 3' phosphoramidite or a compound of formula I where Z is a group of formula II is used, a coupling cycle involves an oxidation or sulphurisation step, while where a 3' H-phosphonate is used oxidation or sulphurisation is effected after chain extension is complete, and where a 3' phosphodiester is used no oxidation is required.


When an oligonucleotide having the desired number of nucleosides has been synthesised, it may be treated to remove hydroxy-protecting groups such as dimethoxytrityl groups, using conventional methods, for example by treatment with an aqueous organic acid such as trifluoroacetic acid. Where the oligonucleotide has been synthesised on a solid support, this treatment may be effected before or after detachment of the oligonucleotide from the support, for example using conventional methods such as treatment with concentrated aqueous ammonia, which treatment also removes a protecting group which may have been present on an exocyclic nitrogen atom in one or more of the nucleosides used in the synthesis of the oligonucleotide. The terminal 5' hydroxyl generated on deprotection can be reacted to introduce a 5' terminal modifying group, such as a phosphate group or other 5' modifying group, for example using the procedures described by Beaucage and Iyer, *Tetrahedron* 49, 1925 - 63 (1993).

The invention is illustrated by the following Examples.


In the formulae of compounds of the Examples and their precursors, Et is ethyl, Me is methyl, T is 1-thyminyl, t-Bu is tert-butyl, Ph is phenyl and DMTr is dimethoxytrityl.

Compounds used in the Examples and precursors thereof are prepared as follows:

Compound A




This is Compound 87 of WO 96/08503, prepared as described in Example 84 of WO 96/08503.


Compound B

To a solution of Compound A (0.85g, 1.22mmol) in anhydrous methanol (10ml) is added sodium methoxide (1.5ml 4.4N solution in methanol). After stirring for 16 hours at room temperature, concentration and purification by flash silica column chromatography (gradient elution - chloroform, methanol, triethylamine 100:20:1-100:35:1), followed by further purification by passing a solution of the product in aqueous 0.5% triethylamine through a Dowex 50W-X2 ion exchange column (triethylamine form) gives, after concentration, Compound B.

^{31}P nmr ^1H decoupled (CD_3OD , 162MHz) δ 23.7 ppm.

Compound C

This is Compound 84 of WO 96/08503, prepared as described in Example 81 of WO 96/08503.

Compound D

To a solution of Compound B (500 mg, 0.71mmol) and dicyclohexylcarbodiimide (189mg, 0.92mmol) in dry THF (5.4ml) under argon at room temperature is added 3-hydroxy propionitrile (58 μ l, 0.85mmol). The resulting solution is heated at 55°C for 2 hours. After cooling the mixture is filtered and diluted with ethyl acetate (20ml) and washed with water and brine, dried over Na₂SO₄, filtered and concentrated. The resulting product is taken up in dichloromethane (5ml) and filtered and concentrated, this process being repeated as required to remove dicyclohexyl urea, to give Compound D, isolated as a mixture of diastereoisomers at phosphorus.


Alternative Method: To solution of Compound B (12.3g, 17.4mmol) and dicyclohexylcarbodiimide (3.94g, 19.1mmol) in THF (120ml) under argon is added 3-hydroxypropionitrile (1.42ml, 20.8mmol) and the resulting mixture is heated to 60°C for 4 hours. Concentration to approximately half volume is followed by dilution with an equivalent volume of ethyl acetate and filtration to remove the resultant precipitate. The organic phase, which after washings of the filtrate with ethyl acetate is now approximately 200ml, is washed with water (4x80ml) and saturated brine (6x80ml), dried over sodium sulphate and concentrated. The resulting solid is redissolved in acetonitrile (200ml) and extracted with hexane (6x50ml). The acetonitrile phase is then concentrated and dried under high vacuum to give Compound D.

Found C 62.2%; H 5.9%; N 6.5%

C₃₅H₃₈N₃O₈P.H₂O requires C 62.05%; H 5.95%; N 6.2%


^{31}P nmr ^1H decoupled (162 MHz, CDCl_3) δ 37.5, 37.3 ppm.

Compound E



This is Compound 100 of WO 96/08503, prepared as described in Example 98 of WO 96/08503.

Compound F



To a solution of Compound E (0.47g, 0.60mmol) dicyclohexylcarbodiimide (DCC) (0.148g, 0.72mmol) and triethylamine hydrochloride (0.24g, 1.8mmol) in anhydrous chloroform (3ml) under argon is added 3-hydroxypropionitrile (81 μl , 1.2mmol). The resulting solution is heated at 60°C for four hours. The mixture is then concentrated under vacuum, taken up in

16


ethyl acetate (50ml) and filtered. The resulting clear solution is extracted with water (3x20ml) and saturated brine (8x20ml). Drying over sodium sulphate, filtration and concentration under reduced pressure gives crude product containing DCC. The DCC is removed by dissolution of this crude product in acetonitrile and repetitive extraction with hexane.

Found C 63.35, H 6.50, N 8.80%;

$C_{40}H_{46}N_5O_7P \cdot H_2O$ requires C 63.40, H 6.40, N 9.25%.


^{31}P nmr 1H decoupled (162 MHz, $CDCl_3$) δ 38.1, 37.4 ppm.

Compound G




To a solution of Compound B (1.0g, 1.41mmol) in dichloromethane (15ml) under an argon atmosphere at 5°C is added triethylamine (98 μ l, 0.71mmol) followed by the dropwise addition of methylchloroformate (142 μ l, 1.84mmol). After 5 minutes the solution is allowed to warm to room temperature, and is stirred for 45 minutes. The mixture is diluted with CH_2Cl_2 and washed with water, dried (Na_2SO_4) and concentrated. Purification by flash silica column chromatography (eluent ethylacetate:methanol 10:1) gives Compound G as a mixture of diastereomers at phosphorus.

^{31}P nmr 1H decoupled ($CDCl_3$, 162 MHz) δ 37.3 and 37.0 ppm. 1H nmr ($CDCl_3$, 500 MHz) δ 8.10 (1H, brs, NH), 7.55 (1H, s, H6), 7.35 (2H, d, ArH), 7.25 (6H, m, ArH), 7.15 (1H, m, ArH), 7.00 (1H, d, PH , J 530Hz), 6.80 (4H, d, ArH), 3.75 (1H, m, H4'), 3.70 (6H, s, $ArOCH_3$), 3.55 (1H, dd, H5'), 3.25 (1H, dd, H5'), 2.70 (1H, m, H3'), 2.45 (1H, m, H2'), 2.20 (1H, m, H2'), 1.75 (1H, m, H6'), 1.60 (1H, m, H6'), 1.45 (3H, brs, TCH_3) ppm.

Compound H

Triazole (20.7g, 0.3mol) is carefully dried by coevaporation with dry pyridine (4x50ml), then dissolved in pyridine (80ml) and cooled to 0°C under argon. Triethylamine (41.8ml, 0.3mol) is added and the mixture stirred during the addition of phosphorus oxychloride (6.87ml, 75 mmol). The resulting solution is stirred at 0°C for 15 minutes. A solution of carefully dried (coevaporation 3x50ml pyridine) Compound 83 of WO 96/08503, prepared as described in Example 80 of WO 96/08503 (20.6g, 30mmol) in pyridine (50ml) is added dropwise. The resulting strongly coloured solution is allowed to warm to room temperature. After 18 hours, the mixture is concentrated and dissolved in THF (300ml). Concentrated ammonia is added (90ml) and the mixture is stirred at room temperature for 18 hours. Concentration and purification by flash silica column chromatography (gradient elution ethyl acetate plus methanol 10%-25%) gives Compound H as a mixture of diastereoisomers at phosphorus. ^{31}P nmr ^1H decoupled (CD_3OD , 162 MHz) δ 50.0 and 49.5 ppm.

18

Compound J


To a solution of Compound H (7.87g, 11.5mmol) in dry THF (110ml) under argon is added triethylamine (6.4ml, 45.9mmol) and the resulting solution cooled to -70°C during the addition of benzoyl chloride (2.00ml, 17.21mmol). The solution is then allowed to warm to room temperature and is stirred overnight. The resulting mixture is partially concentrated, dissolved in ethyl acetate and washed twice with water and then once with brine. The solution is dried over Na_2SO_4 , concentrated and purified by flash silica column chromatography (gradient elution hexane:ethyl acetate 1:5 - pure ethyl acetate) to give Compound J as a mixture of 2 diastereoisomers at phosphorus.

^{31}P nmr ^1H decoupled (CDCl_3 , 162 MHz) δ 47.0 and 46.8 ppm.

Found C 61.85, H 7.25, N 5.10%

$\text{C}_{42}\text{H}_{56}\text{O}_8\text{N}_3\text{PSi} \cdot 3/2 \text{H}_2\text{O}$ requires C 61.75, H 7.30, N 5.15%


19

Compound K

To a solution of Compound J (14.5g, 18.34mmol) in dry THF (160ml) is slowly added a solution of tetrabutylammonium fluoride in THF (20.2ml, 20.2mmol). The resulting solution is stirred at room temperature for 6 hours, evaporated to give a yellow oil and then purified by flash silica column chromatography (gradient elution ethyl acetate - ethyl acetate:ethanol 7:1) to give Compound K as a mixture of 2 diastereoisomers at phosphorus).

^{31}P nmr ^1H decoupled (CDCl_3 , 162 MHz) δ 48.9 and 48.8 ppm.

Found C 55.65, H 7.00, N 7.35% $\text{C}_{26}\text{H}_{38}\text{O}_8\text{N}_3\text{P} \cdot \frac{1}{2} \text{H}_2\text{O}$ requires C 55.70, H 7.00, N 7.50%

Compound L

Compound K (2.48g, 4.50mmol) is carefully dried by coevaporation from dry toluene (3x20ml). Dichloromethane (45ml) is added followed by ethanol (2.62ml, 45mmol), all under argon at room temperature. The resulting solution is stirred and cooled to -15°C during the


20

addition of trimethylsilyl chloride (5.69ml, 45.0mmol). The resulting mixture is thoroughly degassed with argon and slowly warmed to room temperature. After stirring at room temperature for 6 hours the mixture is diluted with dichloromethane and washed twice with sodium bicarbonate solution, dried over Na_2SO_4 and concentrated to give Compound L as a mixture of 2 diastereoisomers at phosphorus.

^{31}P nmr ^1H decoupled (CDCl_3 , 162 MHz) δ 35.5 and 34.2 ppm.

Found C 55.5, H 6.10, N 9.70% $\text{C}_{20}\text{H}_{26}\text{N}_3\text{O}_6\text{P}$ requires C 55.15, H 6.00, N 9.65%


Compound M



Compound L (4.79g, 11.0mmol) is carefully dried by coevaporation with dry pyridine (6x50ml) and then dissolved in pyridine (80ml) before addition of triethylamine (3.1ml, 22mmol) followed by dimethoxytrityl chloride (4.10g, 12.1mmol) at room temperature under argon. After 16 hours at room temperature the mixture is concentrated and stored at -20°C overnight. The resulting solid is dissolved in THF (50ml), triethylamine (15ml) and water (10ml) and stirred at room temperature for 7 hours. The resulting mixture is concentrated and purified by flash silica column chromatography (gradient elution ethyl acetate, triethylamine 50:1 - ethyl acetate, methanol, triethylamine 55:45:2) to give Compound M.

^{31}P nmr ^1H decoupled (CDCl_3 , 202 MHz) δ 22.4 ppm.


Found C 64.5, H 6.65, N 6.4% $\text{C}_{45}\text{H}_{55}\text{N}_4\text{O}_6\text{P} \cdot 3/2\text{H}_2\text{O}$ requires C 64.5, H 6.95, N 6.7%

Compound N


Compound N is prepared by the alternative method described for Compound D, but using Compound M as starting material in place of Compound B, and omitting acetonitrile-hexane partitioning.

^{31}P nmr ^1H decoupled (CDCl_3 , 162 MHz) δ 37.2 and 36.8 ppm.

MS (FAB) m/z 763 (MH^+).

Example 1

Compound 1




Compound 2


To a solution of carefully dried Compound C (40mg, 70μmol) in deuteriochloroform (0.5ml) containing pyridine (80μl, 1mmol) is added dichlorotriphenylphosphorane (113mg, 350μmol). The resulting mixture is shaken to dissolve the phosphorane and then allowed to stand at ambient temperature. The progress of the reaction is monitored by ³¹P nmr. The product, Compound 1, resonates at 201 ppm. After 16 hours, additional dichlorotriphenylphosphorane (45mg, 140μmol) is added. After an additional 24 hours, ³¹P nmr shows the reaction to be 95% complete. A total (150μl, 1.07mmol) of diisopropylamine is added in 30μl portions to the crude reaction mixture at -30°C. The resulting mixture is allowed to warm to room temperature and then diluted with dichloromethane (20ml), washed twice with deionised water (2 x 10ml), dried (Na₂SO₄) and concentrated. Purification by flash silica column chromatography (gradient elution 4:1 - 2:1 hexane-ethylacetate) gives Compound 2 as a mixture of 2 diastereoisomers.

³¹P nmr ¹H decoupled (CDCl₃, 162 MHz) δ 123.0, 122.3.

Example 2



Compound 3




Compound 4


Using the procedure of Example 1, Compound D (30mg, 44μmol) is converted firstly into Compound 3 and then, replacing the diisopropylamine used in Example 1 by pyrrolidine (56μl, 0.67mmol), into Compound 4.

Example 3

Oligonucleotide 1, having the sequence 5'- TTT tTC TCT CTC TCT -3' where t represents a nucleoside unit derived from Compound 4, is prepared by standard solid phase phosphoramidite oligonucleotide synthesis as described in 'Oligonucleotides and Analogues A Practical Approach' ed. F. Eckstein, IRL Press 1991, except that Compound 4 is used instead of the usual 3'-phosphoramidite - substituted nucleoside at the appropriate point in the synthesis, so that a compound of formula I where Z is a group of formula III in which Nuc is a residue of the oligonucleotide TC TCT CTC TCT is formed and then oxidised to an oligonucleotide of formula VIII where X is oxygen in the standard oxidation step, this oligonucleotide being coupled further to give Oligonucleotide 1.

Example 4

Compound 3



Compound 5

To a stirred solution of bis(trichloromethyl)carbonate (1.01g, 3.40mmol) in dry dichloromethane (10ml) under an argon atmosphere at 5°C is added a solution of triphenylphosphine (2.68g, 10.21mmol) in dry dichloromethane (8ml + 7ml washings). The mixture is immediately allowed to warm to room temperature over 5 minutes, pyridine (1.24ml) is then added and the mixture thoroughly mixed and recooled to 5°C. To the resulting pale brown solution is added a solution of Compound D (3.45g, 5.11mmol) in dry dichloromethane (10ml + 5ml washings) containing pyridine (1.24ml). On completion of

24

addition the solution is allowed to warm to room temperature. After stirring at room temperature for 1 hour the mixture contains Compound 3 (^{31}P nmr δ 206.7 ppm). The mixture is cooled to -30°C and diisopropylamine (3.57ml, 27mmol) added over 5 minutes. The mixture is allowed to warm to room temperature over 10 minutes and then concentrated to half volume under vacuum. This solution is immediately transferred onto the top of a flash silica chromatography column (300g, silica) eluting with hexane:tetrahydrofuran:triethylamine 25:25:1. Some pure product fractions result. Impure fractions are subjected to further chromatography on flash silica, eluting with hexane:tetrahydrofuran:dichloromethane:triethylamine 25:25:10:1. All product fractions are pooled and concentrated to give Compound 5 isolated as a white foam and as a mixture of diastereoisomers at phosphorus.


Found C 66.2; H 7.25, N 7.5%; $\text{C}_{41}\text{H}_{51}\text{O}_7\text{N}_4\text{P}$ requires C 66.3, H 6.9, N 7.55%.

^{31}P ^1H decoupled (162MHz, CDCl_3) δ 128.3, 127.8 ppm.


M/Z (ES $^+$) 743 (M+H) $^+$

(ES) 741 (M-H) $^-$.

Example 5



Compound 6



Compound 7


25

Compounds 6 and 7 are prepared by the method of Example 4 but substituting Compound F for Compound D. The reaction is carried out with the same mole quantities of reagents but at half the concentration. In the initial phase of the preparation the reaction is allowed to proceed for 2 hours rather than 1 hour to give Compound 6 (^{31}P nmr δ 205.7 ppm). The final work up involves washing the crude product twice with water, concentration and purification by flash silica column chromatography (eluant ethyl acetate/methanol 10:1 + 2% Et_3N) to give Compound 7 as a mixture of 2 diastereomers.


Found C 63.85, H 7.05, N 9.55% $\text{C}_{46}\text{H}_{59}\text{N}_6\text{O}_6\text{P} \cdot 2\text{H}_2\text{O}$ requires C 64.25, H 7.35, N 9.80%.

^{31}P nmr ^1H decoupled (CDCl_3 , 162MHz) δ 128.6 and 128.1 ppm.

M/Z (ES^+) 823 (MH^+).

Example 6

Compound 9



Compound 8


Compounds 8 and 9 are prepared by the method of Example 4 but substituting Compound G for Compound D. The initial phase of the reactions gives Compound 8 (^{31}P nmr δ 207.7 ppm). Final purification is performed by repetitive flash silica column chromatography, firstly eluting with (hexane/ethyl acetate 2:1 + 2% Et_3N) and subsequently with (hexane/diethyl ether 1:3 + 2% Et_3N) to give Compound 9 as a mixture of 2 diastereomers.

^{31}P nmr ^1H decoupled (CDCl_3 , 202MHz) δ 128.8 and 127.8 ppm.

MZ (ES^+) (704, 705, 706, MH^+)

(ES^-) (702, 703, 704 M-H^-).


26

Example 7


Compound 10

Compound 10 is prepared by the method of Example 6, but replacing the diisopropylamine used in Example 6 by dimethylamine, the intermediate Compound 8 being reacted with a 2.0molar solution of dimethylamine in THF. Final purification by flash silica column chromatography (eluant ethyl acetate/hexane 3:1 + 2% Et₃N) gives Compound 10 as a mixture of 2 diastereomers.

³¹P nmr ¹H decoupled (CDCl₃, 162MHz) δ 144.3 and 144.2 ppm.

Example 8

Compound 11



Compound 12

Compounds 11 and 12 are prepared by the method of Example 10, but substituting Compound N for Compound D. In this case the final product Compound 12 is purified by flash silica column chromatography (hexane-THF-triethylamine 50:50:1).

^{31}P nmr ^1H decoupled (CDCl_3 , 162MHz) δ 128.1 and 127.6 ppm.

MS (ES $^+$) 846, 847, 848 (MH) $^+$

(ES $^-$) 844, 845, 846 (M-H) $^-$

Examples 9-15

The following modified oligonucleotides are prepared by standard DNA synthesis technology using a phosphoramidite strategy. All couplings are performed as double couplings, that is after exposing the support bound oligonucleotide to the phosphoramidite solution or modification solution for a few seconds, the solution is displaced by a fresh batch of monomer solution. In the case of coupling with phosphoramidites, the support is washed before oxidation or thiation (sulphurisation). In the case of coupling with the modified monomers of formula I, where Z is a group of formula II, also known as phosphonamidites, the support is

immediately exposed to the oxidation or thiation solution with no wash cycle. This procedure minimises potential hydrolysis of the support bound PIII intermediate.

In the nucleotide sequences for the modified oligonucleotides given below, T, C, A and G have their usual significance and are derived from the usual phosphoramidite monomers, t denotes a nucleotide unit derived from Compound 5 and c denotes a nucleotide unit derived from Compound 7, Compound 5 and Compound 7 being used at the appropriate points in the syntheses. Sulphurised internucleotide linkages are denoted by s; all other internucleotide linkages are oxidised.

Example 9

5' TTTTtCTCTCTCTCT 3'

TOF MS	Found	=	
	Required	=	4424.3

Example 10

5' ttttCTCTCTCTCT 3'

TOF MS	Found	=	4415.3
	Required	=	4415.9

Example 11

5' GCGttttttttGCG 3'

TOF MS	Found	=	4859.8
	Required	=	4865

Example 12

5' TTTTTTTTTTTTTTTT 3'

TOF MS Found = 5709.3

Required = 5711.7

Example 13

5' ttctcG₃C₃C₃C₃G₃C₃T₃C₃C₃tcctcC 3'

TOF MS Found = 6086.6

Required = 7001.5

Example 14

5' tstscstscsGsCsCsCsGsCsTsCsCstscscstscsC 3'

TOF MS Found = 6214.5

Required = 6246.1

Example 15

5' tstscstscsGsCsTsGsGsTsGsAsGstststscsA 3'


TOF MS Found = 6092.9

Required = 6114


30

Claims

1. A compound of formula



where B^1 is a radical of a nucleoside base, R^1 is hydrogen or a hydroxy-protecting group, R^2 is hydrogen, hydroxy or a 2' nucleoside-modifying atom or group, R^3 is an unsubstituted or substituted C_1 to C_{10} alkyl, C_2 to C_{10} alkenyl, C_4 to C_{10} cycloalkylalkyl, C_6 to C_{10} aryl or C_7 to C_{13} aralkyl group, and Z is halogen or a group of formula



where R^4 and R^5 are each independently an unsubstituted or substituted C_1 to C_{10} alkyl, C_2 to C_{10} alkenyl, C_4 to C_{10} cycloalkylalkyl, C_6 to C_{10} aryl or C_7 to C_{13} aralkyl group, or R^4 is said group and R^5 is hydrogen, or R^4 and R^5 together with the nitrogen atom to which they are attached denote a five- to thirteen- membered heterocyclic ring, or Z is a group of formula




where Nuc is the residue of a natural or synthetic nucleoside or oligonucleotide after removal of a 5'-hydroxyl group therefrom attached through a 5' methylene thereof to the indicated oxygen atom.

2. A compound according to claim 1, in which the substituted groups are substituted by halogen, hydroxy, C_1 - C_4 alkoxy, cyano, nitro or tri(C_1 - C_{15} hydrocarbyl)silyl.

3. A compound according to claim 1 or 2, in which R^1 is hydrogen, a C_7 to C_{40} araliphatic group or a tri(C_1 - C_{15} hydrocarbonyl) silyl group.
4. A compound according to claim 3, in which R^1 is hydrogen, an unsubstituted or substituted triphenylmethyl group or C_1 - C_8 alkyl di(C_6 - C_8 aryl)silyl group.
5. A compound according to claim 4, in which R^1 is a dimethoxytrityl or tert-butyl diphenylsilyl group.
6. A compound according to any one of the preceding claims, in which R^2 is hydrogen, halogen, hydroxy or a group of formula $-OR^6$ or $-OCOR^6$ where R^6 is an unsubstituted or substituted C_1 to C_{10} alkyl, C_2 to C_{10} alkenyl, C_3 to C_8 cycloalkyl, C_4 to C_{10} cycloalkylalkyl, C_6 to C_{10} aryl or C_7 to C_{25} aralkyl group or a tri(C_1 - C_{15} hydrocarbonyl)silyl group.
7. A compound according to claim 6, in which R^2 is hydrogen, hydroxy, fluorine, a group of formula $-OR^6$ where R^6 is a C_1 to C_4 alkyl group, a C_2 to C_4 alkenyl group, a C_7 to C_{13} aralkyl group, or a group of formula $-(R^7-O)_n-R^8$ where R^7 is C_1 to C_4 alkylene, R^8 is C_1 to C_4 alkyl and n is 1 to 8; or R^2 is a group of formula $-OCOR^6$ where R^6 is C_1 to C_4 alkyl or C_6 to C_8 aryl.
8. A compound according to claim 6, in which R^2 is hydrogen, hydroxy, fluorine, methoxy, ethoxy, allyloxy, benzyloxy, methoxyethoxy, $-O-(CH_2-CH_2-O)_7-CH_3$, acetoxy or benzoyloxy.
9. A compound according to any one of the preceding claims, in which R^3 is an unsubstituted or substituted C_1 to C_4 alkyl, C_3 to C_5 alkenyl, C_6 to C_8 cycloalkylalkyl or C_7 to C_9 aralkyl group.


10. A compound according to claim 9, in which R^3 is methyl, ethyl, 2-cyanoethyl, 2,2,2-trichloroethyl, allyl, benzyl or 2-(trimethylsilyl)ethyl.
11. A compound according to any one of the preceding claims, in which Z is a group of formula II in which R^4 and R^5 are independently an unsubstituted or substituted C_1 to C_4 alkyl, C_3 to C_5 alkenyl, C_6 to C_8 cycloalkylalkyl or C_7 to C_9 aralkyl group, or R^4 is C_1 to C_4 alkyl or C_7 to C_9 aralkyl and R^5 is hydrogen, or R^4 and R^5 together with the attached nitrogen atom denote a five- or six- membered heterocyclic ring having one or two heteroatoms in the ring.
12. A compound according to claim 11, in which R^4 and R^5 are independently methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, allyl or benzyl or R^4 and R^5 together with the attached nitrogen atom denote a pyrrolidine, piperidine or morpholine ring.
13. A compound according to any of claims 1 to 10, in which Z is a group of formula III in which Nuc is attached to a solid support.
14. A compound according to claim 1, in which Z is chlorine or bromine.
15. A compound according to any of the preceding claims, which is of formula



where B^1 , R^1 , R^2 , R^3 and Z are as defined in any of the preceding claims.

16. A compound according to claim 1 or claim 15 in which B¹ is thyminy, 5-methylcytosiny, or thyminy or 5-methylcytosiny substituted on a nitrogen atom by a protecting group; R¹ is dimethoxytrityl or tert-butyldiphenylsilyl; R² is hydrogen; R³ is methyl, ethyl or 2-cyanoethyl; and Z is chlorine or a group of formula II where R⁴ and R⁵ are each methyl or isopropyl or Z is a group of formula III.


17. A method of preparing a compound according to claim 1 in which Z is a group of formula II which comprises reacting a compound of formula I in which Z is halogen with a compound of formula



where R⁴ and R⁵ are as defined in claim 1.

18. A method according to claim 17, in which the reaction is carried out in an organic solvent in the presence of a tertiary nitrogen base at a temperature from -78°C to 50°C.

19. A method of preparing a compound of formula I in which Z is halogen which comprises non-oxidatively halogenating a compound of formula




where B¹, R¹, R² and R³ are as defined in claim 1 with a non-oxidative halogenating agent.

20. A method according to claim 19, in which the non-oxidative halogenating agent is a halophosphorane.

21. A method according to claim 19 or 20, in which the halogenation is carried out in the presence of a base in an organic solvent at a temperature from -20°C to 60°C.


22. A method of preparing a compound according to claim 1 in which Z is a group of formula III which comprises reacting a compound according to claim 1 where R¹ is a hydroxy-protecting group and Z is a group of formula II with a natural or synthetic nucleoside or oligonucleotide having a free 5' hydroxyl group.

23. A method of preparing an oligonucleotide of formula




where B¹, R¹, R³ and Nuc are as defined in claim 1 and X is oxygen or sulphur, which comprises subjecting a compound according to claim 1 where Z is a group of formula III to oxidation to give an oligonucleotide of formula VIII where X is oxygen or to sulphurisation to give an oligonucleotide of formula VIII where X is sulphur.

24. A method of preparing an oligonucleotide which comprises subjecting a nucleoside having a protected 5' hydroxy group and, at the 3' position, a group of formula




where R³, R⁴ and R⁵ are as defined in claim 1, to a nucleoside coupling reaction with a natural or synthetic nucleoside or oligonucleotide having a free 5' hydroxy group, to form an oligonucleotide precursor having an internucleoside linkage of formula

35



where R^3 is as defined in claim 1, and converting the precursor into an oligonucleotide having an internucleoside linkage of formula



where R^3 is as hereinbefore defined and X is oxygen or sulphur by oxidising the precursor to give an oligonucleotide having an internucleoside linkage of formula XI where X is oxygen or sulphurising the precursor to give an oligonucleotide having an internucleoside linkage of formula XI where X is sulphur.

25. Use of a compound according to claim 1 in the synthesis of an oligonucleotide.
26. A compound according to claim 1 substantially as described in any of the Examples.
27. A method according to claim 17, substantially as described in any of Examples 1, 2 or 4 to 8.
28. A method according to claim 19, substantially as described in any of Examples 1, 2, 4, 5, 6 or 8.
29. A method according to claim 22, 23 or 24, substantially as described in any of Examples 3 and 9 to 15.